

Technical University of Denmark



Environmental radioactivity during 50 years

Nielsen, Sven Poul

Publication date:
2013

[Link back to DTU Orbit](#)

Citation (APA):

Nielsen, S. P. (2013). Environmental radioactivity during 50 years [Sound/Visual production (digital)]. NKS Workshop on Radioanalytical Chemistry, Roskilde, Denmark, 02/09/2013, http://www.nks.org/en/seminars/presentations/nks-b_radioanalysis_workshop_2-6_september_2013.htm

DTU Library

Technical Information Center of Denmark

General rights


Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Environmental radioactivity during 50 years

Sven P. Nielsen






Investigations of man-made radioactivity in the Danish environment from 1957




Atmospheric nuclear weapons tests

Global pollution from atmospheric nuclear weapons tests: fission products, activation products, fissile material and tritium




Radioecological sensitivity



- Studies covering
 - Air, water, soil
 - Grain, bread
 - Grass
 - Vegetables and fruit
 - Sea plants
 - Milk, meat, fish
 - Total diet
 - Human body, bone
- Radioecological sensitivity is the time integral of quantities of the sample type from a quantity of the radionuclide deposited
- Example for Cs-137 in Danish cow's milk
 - 2.0 Bq/L d per Bq/m²

Strontium-90 and caesium-137

- Fission products of particular importance due to long half lives and significant uptake in food chains



Aerosols


- Monitoring of radioactivity in air is based on aerosol collectors located in Haderslev, Allinge and Risø.
- Air is sampled at flow rates of 500-2000 m³/h through organic filters retaining particles
- Filters are changed weekly and analysed for short-lived radionuclides first and later for longer lived radionuclides, particularly ⁷Be, ²¹⁰Pb, ⁹⁰Sr, ¹³⁷Cs.



Risø




Radioactivity in Air at Risø



Precipitation


Precipitation is collected at Risø and 10 other locations in Denmark and analysed for content of Sr-90 and Cs-137




Caesium-137 in precipitation at Risø, 1960-2009



Milk, potatoes, vegetables and total diet



Sampling zones (I-VIII) for milk, potatoes, vegetables and total diet in Denmark.




Caesium-137 in milk collected in 1962-2008.



Caesium-137 in beef and pork collected in 1963-2009

Cereals: rye, oats, wheat, barley






Strontium-90 in rye collected in 1959-2008 and in rye bread collected in 1962-1990.





Grain sampling locations (State experimental farms in Denmark)

Stream, lake and ground water




Sea water and plants




*Caesium-137 in seaweed (*Fucus vesiculosus* and *Fucus seratus*) from February 1983 to June 2009 collected at Klint, Zealand (55°58'N, 11°35'E).*

DTU Nutech, Technical University of Denmark




Caesium-137 in fish/cod




Humans

- Employees at Risø monitored for radiocaesium and tritium
- Human bone samples received from hospitals (with difficulty)




Radiocaesium ($Bq\ ^{137}Cs\ (kg\ K)^{-1}$) in diet, urine, and humans from Zealand in 1986-2009.



Strontium-90 levels in bone from adults (> 29 years) 1960-2009.

External exposure



External exposure rates in 4 locations in Denmark, as measured with a NaI(Tl) detector.




Grain sampling locations (State experimental farms in Denmark)



Radioactive contamination in Denmark

- Poster in building 204 shows concentrations of strontium-90 and caesium-137 in air, precipitation, milk and grass at Risø and in Denmark since the 1950's
- Including input from the Fukushima accident in Japan in 2011



Why monitor environmental radioactivity?

- EURATOM Treaty: Health and safety matters - Obligation of EC Member States to monitor levels of radioactivity in air, soil and water and to ensure compliance with basic standards
- Helsinki Convention: Contracting Parties undertake to prevent and eliminate pollution of the marine environment of the Baltic Sea Area caused by harmful substances from all sources
- Study man-made and naturally occurring radionuclides in the environment to document baseline levels and increase knowledge on behaviour and processes
- Expertise available for emergency purposes in case of accidents/incidents involving release of radioactivity to environment
- Improve radiological assessment models in decision support systems used by authorities in case of accidents
- Useful platform for research and development of existing and new analytical methods and application of these in other areas